

Assessment & Management of Psychological and Sexual Issues in LGBTQ+ populations

Avinash De Sousa¹

Amrit Patojoshi²

Navendu Gaur³

¹Consultant Psychiatrist and Founder Trustee, Desousa Foundation, Mumbai.

²Professor and Head, PG Department of Psychiatry, Hi-Tech Medical College and Hospital, Bhubaneswar.

³Consultant Psychiatrist and Director, Gaur Mental Health Clinic, Ajmer.

Address for correspondence –

Dr Avinash Desousa, Carmel, 18, St. Francis Road, Off SV Road, Santacruz West, Mumbai 400054. Tel - +91-9820696828. E-mail – avinashdes888@gmail.com

Assessment & Management of Psychological and Sexual Issues in LGBTQ+ populations

Executive Summary

LGBTQ+ individuals experience disparities in psychological health, arising not from sexual orientation or gender diversity themselves, but from minority stress, stigma, discrimination, victimization, family rejection, violence, and barriers to equitable healthcare. This clinical guideline provides a framework for assessment and management of psychological and sexual concerns, emphasizing gender-affirmative, sexuality-affirmative, confidential, and trauma-informed care. Sexual concerns include desire, arousal, orgasm, erectile function, genital pain, sexual satisfaction, body image, and safer-sex practices. Assessment should include sexual, psychiatric, medical, psychosocial, substance-use, relationship, family, trauma, and minority-stress histories, while avoiding assumptions about identity, relationships, or sexual practices. The guideline highlights limited epidemiological evidence in India, particularly for lesbian, bisexual, transmasculine, and queer populations, despite documented vulnerabilities among MSM and transgender populations. Structured assessment of depression, anxiety, suicidality, self-harm, PTSD, substance use, gender dysphoria, family support, discrimination, and social connectedness is recommended. Management involves treatment of medical and hormonal contributors, HIV/STI prevention, psychotherapies including CBT, DBT, IPT, EMDR, mindfulness-based therapy, and ACT, and pharmacotherapy when indicated. Gender-affirming psychotherapy supports identity exploration, coping, trauma recovery, and resilience, while family psychoeducation and community support may strengthen acceptance. Conversion or reparative therapy is explicitly discouraged. The guideline advocates shared decision-making, informed consent, fertility counselling before gender-affirming interventions, affirming documentation, confidentiality, and multidisciplinary collaboration involving psychiatrists, psychologists, sex therapists, endocrinologists, surgeons, social workers, speech therapists, occupational therapists, peer-support workers, and legal professionals. Regular follow-up is essential to monitor mental health, treatment response, transition-related care, sexual health, and quality of life. Further research is needed to strengthen the evidence base.

Keywords: sexual issues, sexual, psychosexual, LGBTQ+.

Introduction

Lesbian, Gay, Bisexual, Transgender, Queer, and other gender and sexual minority (LGBTQ+) individuals experience significant disparities in mental and sexual health, largely driven by minority stress, stigma, discrimination, victimization, and barriers to accessing equitable healthcare. Although sexual orientation and gender diversity are normal variations of human sexuality and are not psychiatric disorders, LGBTQ+ individuals are disproportionately affected by depression, anxiety, substance use disorders, suicidality, trauma-related disorders, and sexual health concerns. Despite growing evidence supporting LGBTQ+-affirmative care, many healthcare professionals report limited training and confidence in conducting culturally competent psychological and sexual health assessments. Evidence-based clinical practice guidelines are therefore essential to standardize assessment, promote affirmative and trauma-

informed care, facilitate early identification and management of psychological and sexual health concerns, discourage harmful practices such as conversion therapy, and improve the quality, safety, and equity of healthcare for LGBTQ+ individuals across diverse clinical settings.^[1]

Common sexual dysfunctions in LGBTQ+ individuals include erectile dysfunction in gay, bisexual, and transgender men, hypoactive sexual desire disorder; female sexual arousal disorder in transgender men and lesbian or bisexual women; genital pain disorder, including dyspareunia and vaginismus in lesbian, bisexual, and transgender women; and persistent genital arousal disorder. Psychological problems like depression and anxiety can precipitate or exacerbate these sexual issues. The social stressors that this population encounters are unique and include stigma, discrimination, violence, poor social acceptance, etc. Biological factors like hormonal changes associated with gender-affirming therapies also can contribute to their sexual dysfunction.^[2]

Epidemiology of sexual problems in LGBTQ+ populations in India

The epidemiology of sexual problems among LGBTQ+ populations in India remains incompletely characterized because of stigma, under-reporting, limited population-based research, and the historical focus of Indian sexual-health studies on HIV and sexually transmitted infections (STIs) rather than broader sexual dysfunction. Available evidence indicates substantial sexual-health vulnerabilities among men who have sex with men (MSM) and transgender people, including HIV, syphilis, gonorrhoea, chlamydia, genital herpes, and other STIs. A multicentre study of 4,597 MSM in southern India reported HIV prevalence ranging from 7.6% among panthis to 18.1% among hijra/transgender participants, with inconsistent condom use particularly common in non-commercial partnerships.^[3]

More recent research from Chennai and Mumbai found bacterial STI prevalence of 37.8% and 27.6%, respectively, among MSM, demonstrating substantial ongoing sexual-health morbidity. Sexual problems may also include difficulties with sexual desire, arousal, orgasm, sexual pain, body-image concerns, gender dysphoria-related distress, and difficulties negotiating safer sex. However, robust prevalence estimates for these conditions among lesbian, bisexual, transgender and queer populations are scarce. A national scoping review found that research has disproportionately focused on gay/MSM and transgender women, while lesbian, bisexual and transmasculine populations remain considerably understudied.^[4]

Assessment of psychological and sexual problems

A non-judgmental, safe, clinical environment must be provided to the client, ensuring privacy and confidentiality. The individual must be addressed using his/her affirmed name and pronouns used appropriately and consistently. The gender identity, sexual orientation, and the sex assigned at birth must be assessed politely. The prejudices of the clinician regarding the sexual identity, relationships, and practices should not colour the interview. Any embarrassment on the part of the patient must be addressed with empathy (Table 1 and 2). Their sexual problems must be evaluated, in the background of the psychological, relationship, and social issues faced by them (Table 3).^[5]

Sexual problems

Sexual history must be assessed in detail. In homosexual relationships, the power dynamics of the partners are different from that of heterosexual ones and can be reversed. The ‘top’ or

‘bottom’ positions can be reversed. Non-coital sex, like oral sex may be practised. Penetration need not take place. A gay man can hide his sexual dysfunction, like erectile dysfunction (ED), by taking the receptive role in penetrative sex (Table 3).^[5]

To assess sexual dysfunction in a gay man, questions related to libido or interest, sexual arousal, performance, and ejaculation/orgasm/satisfaction must be asked. For a lesbian, in addition to questions on interest or libido, and arousal, those related to orgasm/satisfaction, and dyspareunia are needed. Premature ejaculation (PE) can be seen in men who have sex with men (MSM). Questionnaires like the Arizona Sexual Experiences Scale, the Brief Male Sexual Function Inventory, and the Derogatis Interview for Sexual Functioning – Self Report are some of the questionnaires which can be used for assessing sexual dysfunction, but they have not been validated in this population.^[6]

Psychosocial problems

Psychological factors have an important role in maintaining good sexual function, especially in those with different sexual orientation. The factors which predispose, precipitate, and maintain sexual dysfunction should be assessed in detail. Restrictive upbringing, early traumatic sexual experiences, troubled family relationships, exposure to domestic violence in childhood, etc. can predispose a person to sexual dysfunction. Relationship issues with current partner, infidelity, lack of intimacy, etc. can be precipitating factors. Performance anxiety, lack of proper communication between partners, etc. can maintain the dysfunction. Psychological problems like depression, anxiety, poor self-esteem, guilt about the relationship, etc. must be looked for and evaluated in detail. Interpersonal problems between the partners, discrepancies in the expectations about the partner, etc. must be assessed. Social factors can contribute to the sexual dysfunction, and a detailed assessment is needed. Minority stress, internalized homophobia or bi-negativity can affect sexual satisfaction and functioning. The stigma, lack of acceptance by the family, discrimination, and bullying from society can be internalized, leading to low self-esteem, fear of identifying one’s identity in public, and poor emotional health (Table 4, 5 and 6).^[7]

Substance use

History of substance use to enhance sexual experiences, and the pattern of substance use – whether it amounts to substance use disorder, must be assessed. It must be evaluated whether the substance use is causing sexual dysfunction or other psychological problems. Risk of HIV, HBV infections, etc., associated with IV drug abuse must be assessed.^[8]

Physical illnesses and treatment history

Any medical illnesses, especially chronic ones, like diabetes mellitus or hypothyroidism, that can affect sexual functioning have to be looked for. Medications taken by the patient, including over-the-counter ones, must be evaluated. Those that can contribute to sexual dysfunction have to be identified and deprescription attempted, if possible. The impact of gender affirmation surgeries and medications, including hormonal therapy, on sexual function must be investigated. Evaluation and counselling for sexually transmitted infections, like HIV, must be undertaken.^[9]

Physical examination

Detailed physical examination is needed with due consideration to the privacy and confidentiality of the patient.

Investigations

Blood investigations including blood sugar, fasting lipid profile, hormonal levels, serological test for syphilis, and ELISA for HIV are warranted. For dyspareunia, microscopy and culture may be done as required.

Management (Table 7 to Table 12)

If medical illnesses contributing to sexual dysfunction are identified, they are to be addressed effectively. Surgical procedures and hormonal therapy may be required. Referral to other specialists, like the plastic surgeon or endocrinologists, for medical or surgical interventions, can be helpful. Psychosocial issues must be addressed. Gender-affirmative and sexuality-affirmative care must be provided. It is ideal to work with a multidisciplinary team.

For hypoactive sexual desire disorder, psychological problems like depression or anxiety, societal pressures and relationship-related stressors must be addressed. Hormonal factors also are to be considered and addressed. Female sexual arousal disorder may also be related to psychological factors like gender dysphoria, past trauma, or hormonal problems due to gender-affirming medical treatment. Social factors and interpersonal problems related to relationships also must be addressed.^[10]

In lesbian, bisexual or transgender women, dyspareunia or vaginismus can occur. may be the result of anatomical differences following gender-affirmation surgeries, hormonal changes, or other psychological problems. In transmen, without a history of bottom surgery, genital pain can be the consequence of testosterone therapy or physical dysphoria. Trauma-informed care, counselling, and pelvic floor therapy are needed for proper management of pain.^[11]

Persistent genital arousal disorder is unprovoked, not linked to sexual desire, and distressing. A multidisciplinary approach involving psychiatrists, pelvic floor specialists and neurologists can provide holistic care.

Comorbid psychiatric problems like depression and anxiety must be addressed. Evidence-based psychotherapeutic strategies, like cognitive behaviour therapy (CBT), dialectical behaviour therapy (DBT), eye movement desensitization and reprocessing (EMDR), mindfulness-based therapy (MBT) and acceptance and commitment therapy (ACT) can be provided when indicated. Internalized stigma and rejection sensitivity must be addressed therapeutically. Pharmacotherapy can be provided according to the indication. Risk of suicide must be evaluated and addressed based on an individualized plan. Psychoeducation must be given regarding safer practices while using substances to enhance sexual pleasure, so that informed decision-making is possible.^[12]

Gender affirming psychotherapy using non-judgmental counselling approach can be helpful in exploring gender identity, addressing minority stress and trauma, building coping strategies, and improving mental health and well-being. Sexuality-affirmative therapy validates and advocates for their identities. Their gender identity and sexual orientation are validated, and they are helped to address the emotional trauma of homophobia, biphobia, or transphobia. Conversion or reparative therapy is never recommended. They help families/parents to accept their child's gender and be more supportive. Family psychoeducation and family therapy is provided when needed. Cultural, religious, and family contexts are respected without

stereotyping. Supportive networks and community resources to affirm gender identity and sexuality can help to enhance sexual health and gender affirmation.^[13] Regular follow-up must be ensured to monitor mental health, response to treatment, transition-related care, and quality of life. Evidence for management of sexual dysfunction in LGBTQ+ is not robust. There is a need for more research addressing this area in this population.

Take home points – (Figure 1)

1. Use the individual's affirmed name and pronouns consistently.
2. Create a safe, non-judgmental, confidential clinical environment.
3. Assess gender identity, sexual orientation, and sex assigned at birth respectfully.
4. Avoid assumptions regarding identity, relationships, or sexual practices.
5. Evaluate presenting psychological, sexual, and relationship concerns comprehensively.
6. Screen routinely for depression, anxiety, suicidality, self-harm, and substance use.
7. Assess minority stress, discrimination, bullying, and stigma.
8. Explore family acceptance, rejection, and social support.
9. Screen for childhood trauma, intimate partner violence, and hate crimes.
10. Assess gender dysphoria, transition goals, and expectations.
11. Evaluate readiness and capacity for informed decision-making.
12. Assess sexual function, satisfaction, consent, and sexual health.
13. Provide individualized HIV/STI risk assessment and prevention counselling.
14. Differentiate consensual sexual diversity from psychopathology.
15. Use trauma-informed care throughout assessment and treatment.
16. Employ gender-affirmative and sexuality-affirmative care.
17. Treat psychiatric comorbidities alongside gender-affirming care.
18. Use evidence-based psychotherapies such as CBT, DBT, IPT, EMDR, MBT, and ACT when indicated.
19. Prescribe pharmacotherapy according to standard psychiatric indications.
20. Develop individualized safety plans for suicide or self-harm risk.
21. Address internalized stigma and rejection sensitivity therapeutically.
22. Promote resilience, identity affirmation, and community connectedness.
23. Offer family psychoeducation and family therapy when appropriate.
24. Respect cultural, religious, and family contexts without stereotyping.
25. Discuss fertility preservation before gender-affirming medical interventions.
26. Refer appropriately to endocrinologists, surgeons, sex therapists, speech therapists, and other specialists.
27. Work collaboratively within a multidisciplinary team.
28. Maintain affirming documentation and protect confidentiality.
29. Never recommend conversion or reparative therapy.
30. Ensure regular follow-up to monitor mental health, treatment response, transition-related care, and quality of life.

Table 1. Initial Clinical Assessment

Domain	Assessment Areas	Clinical Questions	Recommended Tools	Red Flags
Demographics	Name, pronouns, gender identity, sex assigned at birth, sexual orientation	"How would you like to be addressed?"	Intake form	Misgendering, assumptions
Presenting Complaint	Psychological, sexual, relational concerns	Duration, severity, triggers	Clinical interview	Suicidality, violence
Gender Identity	Identity development, dysphoria, transition status	Social, legal, medical transition	Gender history interview	Severe gender dysphoria
Sexual Orientation	Identity, attraction, behavior	Outness, identity acceptance	Sexual history	Internalized stigma
Family Support	Acceptance, rejection, abuse	Family relationships	Family assessment	Homelessness, violence
Social Support	Friends, community, workplace	Community connectedness	Social network mapping	Social isolation
Minority Stress	Discrimination, bullying, stigma	Frequency and impact	Minority Stress Scale	Chronic victimization
Trauma	Childhood abuse, hate crimes, IPV	Physical, emotional, sexual trauma	Trauma screening	PTSD symptoms
Psychiatric History	Previous disorders, treatment	Past diagnosis and medications	DSM-5 interview	Untreated severe illness
Substance Use	Alcohol, Nicotine, Other drugs	Pattern, function and dependence	ASSIST, AUDIT	Dependence and overdose risk

224
225
226
227
228
229
230
231
232

Table 2. Mental Health Assessment				
Disorder	Common Presentation Factors	Risk	Assessment Tools	Management
Depression	Low mood, hopelessness	Minority stress, rejection	PHQ-9	CBT, IPT, antidepressants
Anxiety	Worry, panic, avoidance	Bullying, discrimination	GAD-7	CBT, SSRIs
Suicidality	Ideation, attempts	Family rejection,	C-SSRS	Safety planning
Self-harm	Cutting, burning	Emotional dysregulation	Clinical interview	DBT
PTSD	Flashbacks, avoidance	Hate crimes, abuse	PCL-5	Trauma-focused CBT, EMDR
OCD	Obsessions, compulsions	Independent of identity	Y-BOCS	ERP + SSRIs
Eating Disorders	Restriction, bingeing	Body dissatisfaction	EDE-Q	Multidisciplinary care
Substance Use	Coping through substances	Stigma, trauma	ASSIST	MI + relapse prevention
Personality Disorders	Emotional instability	Childhood trauma	SCID-5-PD	DBT, MBT

233
234
235

Table 3. Sexual Assessment

Area	Assessment Components	Clinical Considerations	Intervention
Sexual Function	Desire, arousal, orgasm, pain	Hormones, medications	PLISSIT model
Sexual Satisfaction	Relationship satisfaction	Minority stress	Couples therapy
Sexual Practices	Partners, practices, consent	Avoid assumptions	Psychoeducation
HIV/STI Risk	Condom use, PrEP, testing	Individualized assessment	Prevention counselling
Sexual Compulsivity	Frequency, distress	Rule out mania/OCD	CBT
Fetishes/Kinks	Consensual/non-consensual	Differentiate pathology	Affirmative approach
Consent	Capacity, coercion	IPV screening	Safety interventions
Sexual Trauma	Abuse history	Trauma-informed care	Trauma therapy

236
237
238
239

240
241
242

243
244
245
246
247
248
249
250
251
252

253
254
255
256
257
258
259

Table 4. Gender Dysphoria Assessment

Domain	Assessment	Clinical Indicators	Management
Dysphoria	Persistence	Distress regarding incongruence	Gender-affirmative care
Social Transition	Name, pronouns, clothing	Functioning	Support transition
Medical Transition	Hormones, surgery	Eligibility	Refer to specialists
Expectations	Realistic outcomes	Body image	Psychoeducation
Capacity	Decision-making ability	Informed consent	Shared decision-making
Comorbidity	Psychiatric illness	Stabilization	Parallel treatment

Table 5. Minority Stress Assessment

Stressor	Assessment	Psychological Impact	Clinical Response
Internalized Homophobia	Self-stigma	Depression	CBT
Internalized Transphobia	Shame	Anxiety	Identity affirmation
Rejection Sensitivity	Hypervigilance	Social anxiety	CBT
Concealment	Outness	Stress	Explore readiness
Discrimination	Workplace, healthcare	PTSD	Advocacy
Violence	Hate crimes	Trauma	Trauma therapy
Microaggressions	Daily experiences	Emotional exhaustion	Validation

Table 6. Family Assessment

Domain	Assessment	Protective Factors	Intervention
Acceptance	Level of support	High acceptance	Family therapy
Rejection	Emotional/physical	None	Crisis intervention
Communication	Conflict patterns	Open dialogue	Psychoeducation
Religious Beliefs	Supportive vs rejecting	Inclusive beliefs	Respectful discussion
Financial Support	Dependence	Independence	Social services

Problem	First-line Therapy	Second-line	Pharmacotherapy	Follow-up
Depression	CBT/IPT	ACT	SSRIs	2–4 weeks
Anxiety	CBT	ACT	SSRIs/SNRIs	Monthly
PTSD	TF-CBT	EMDR	SSRIs	Monthly
Self-harm	DBT	MBT	As indicated	Weekly
Substance Use	MI	CBT	Naltrexone etc.	Weekly
Relationship Issues	Couples therapy	EFT	NA	Monthly

261
262
263
264
265
266

Table 8. Gender-Affirmative Care

Intervention	Indications	Professionals	Monitoring
Social Transition	Persistent dysphoria	Mental health team	Adjustment
Voice Therapy	Voice incongruence	Speech therapist	Progress
Hormone Therapy	Informed consent	Endocrinologist	Labs every 3–6 months
Surgery	Persistent dysphoria	Multidisciplinary team	Surgical follow-up
Fertility Counselling	Before hormones/surgery	Reproductive specialist	Documentation

267
268
269
270

Table 9. Sexual Health Management

Issue	Assessment	Intervention	Referral
Erectile Dysfunction	Organic + psychological	CBT, PDE5 inhibitors	Urologist
Low Desire	Hormonal, relational	Sex therapy	Sexologist
Dyspareunia	Medical evaluation	Pelvic therapy	Gynecologist
Orgasmic Disorders	Medication review	Behavioural therapy	Sex therapist
STI Prevention	Risk assessment	PrEP, condoms	Infectious disease specialist

271

272
273
274
275

Table 10. Suicide Risk Assessment

Risk Factor	Low Risk	Moderate Risk	High Risk	Action
Ideation	Passive	Active without plan	Active with plan	Hospitalization if indicated
Previous Attempts	None	One	Multiple	Intensive monitoring
Family Support	Strong	Variable	Absent	Crisis intervention
Substance Use	None	Episodic	Severe	Dual diagnosis treatment
Access to Means	Limited	Possible	Immediate	Means restriction

276
277
278

Table 11. Ethical & Legal Considerations

Area	Best Practice	Avoid
Pronouns	Use affirmed pronouns	Deadnaming
Confidentiality	Protect identity	Unnecessary disclosure
Documentation	Affirming language	Pathologizing language
Conversion Therapy	Never recommend	Any reparative practice
Consent	Shared decision-making	Coercion
Cultural Sensitivity	Respect diversity	Stereotyping

279
280
281
282

Table 12. Multidisciplinary Team Approach

Professional	Primary Role	Referral Indications
Psychiatrist	Diagnosis, medication	Mental illness
Clinical Psychologist	Psychotherapy	Psychological distress
Sex Therapist	Sexual dysfunction	Sexual concerns
Endocrinologist	Hormone therapy	Gender-affirming hormones
Surgeon	Gender-affirming surgery	Surgical transition
Social Worker	Resources, advocacy	Housing, finances
Speech Therapist	Voice training	Voice incongruence
Occupational Therapist	Functional rehabilitation	Disability support
Peer Support Worker	Community support	Isolation
Legal Advisor	Documentation	Legal gender recognition

283
284
285
286

Table 13 Key Principles of Care

Principle	Clinical Recommendation
Respect	Use affirmed name and pronouns consistently
Affirmation	Employ a gender-affirmative and sexuality-affirmative approach
Individualization	Avoid assumptions regarding identity, relationships, or sexual practices
Trauma-Informed Care	Assume trauma may be present and provide safe, non-judgmental care
Cultural Competence	Recognize cultural, religious, and family influences
Evidence-Based Practice	Follow current professional guidelines and standards of care
Collaborative Care	Integrate psychiatry, psychology, endocrinology, primary care, and community support

Prevention	Screen routinely for depression, anxiety, substance use, suicidality, intimate partner violence, and HIV/STIs
Human Rights	Promote dignity, autonomy, confidentiality, and equitable access to healthcare
Follow-Up	Monitor mental health, treatment response, transition-related care, and overall quality of life regularly

Figure 1 - ASSESSMENT FLOW CHART

Presenting Sexual Complaint



Detailed Sexual History



Assess

- Desire
- Arousal
- Orgasm
- Pain
- Satisfaction
- Sexual Practices



Identify Contributing Factors

- Psychological
- Relationship
- Minority Stress
- Medical Disorders
- Hormonal Factors
- Substance Use



Diagnosis



Clinical Diagnosis



Treat Medical Conditions



Address Psychological Problems



Gender & Sexuality Affirmative Care



Psychotherapy
(CBT/DBT/IPT/ACT/EMDR/MBT)



Pharmacotherapy (when indicated)



Family Therapy

Community Support



REFERENCES

1. Wandrekar JR, Nigudkar AS. What do we know about LGBTQIA+ mental health in India? A review of research from 2009 to 2019. *J Psychosexual Health* 2020;2(1):26-36.
2. Sahoo S, Venkatesan V, Chakravarty R. 'Coming out'/self-disclosure in LGBTQ+ adolescents and youth: International and Indian scenario-A narrative review of published studies in the last decade (2012-2022). *Indian J Psychiatry* 2023;65(10):1012-24.
3. Safren SA, Blashill AJ, Lee JS. Sexual orientation and sexual health among men who have sex with men in India. *J Sex Med* 2010;7(7):2346-55.
4. Solomon SS, Mehta SH, Srikrishnan AK, Vasudevan CK, McFall AM, Balakrishnan P, et al. High HIV prevalence and associated risk factors among men who have sex with men in India. *AIDS Behav* 2015;19(11):2077-88.
5. Sathyanarayana Rao TS, Tandon A, Manohar S, Mathur S. Clinical Practice Guidelines for management of Sexual Disorders in Elderly. *Indian J Psychiatry* 2018;60:397-409.
6. Rao TSS, Banerjee D, Tandon A, Sawant NS, Jha A, Manohar S, Rao SS. Psychosexual Health and Sexual Medicine in Consultation–Liaison Psychiatry. *Indian J Psychiatry* 2022;64(Suppl 2):S429–48.
7. Saraff S, Singh T, Kaur H, Biswal R. Stigma and health of Indian LGBT population: A systematic review. *Stigma Health* 2022;7(2):178-95.
8. Chaney MP, Urhahn-Schmitt N. LGBTQ+ substance use and sexual health and wellbeing: A special commentary. *J Counsel Sex Sexual Wellness Res Pract Educ* 2022;4(2):66-9.
9. Pal S, Pathak PK, Rahaman M. Physical-mental health and awareness of HIV/STIs among men who have sex with men in selected cities of India. *BMC Pub Health* 2023;23(1):320.
10. Agrawal N, Shainy MR, Sebastian J, Srivastava A, Scoats R. Sex without Sexual Attraction?: Perceived Sexual Experiences of Indian Sex-favorable Asexuals. *Psychol Sexuality* 2025;16(4):1260-80.
11. Sharan P, Purnima S, Rao R, Kedia S, Khoury B, Reed GM. Field testing of ICD-11 proposals for female sexual dysfunctions in India: Cognitive interviews with patients. *Arch Med Res* 2019;50(8):567-76.
12. Hebbar NY, Jotibannand DY, Kalasagond MB, Hegde SG. Psychiatric morbidities among sexual minorities in India: A cross-sectional study from a north-east state of Manipur. *Natl J Physiol Pharmacy Pharmacol* 2024;14(5):921-4.
13. Gaur PS, Saha S, Goel A, Ovseiko P, Aggarwal S, Agarwal V, Haq AU, Danda D, Hartle A, Sandhu NK, Gupta L. Mental healthcare for young and adolescent LGBTQ+ individuals in the Indian subcontinent. *Front Psychol* 2023;14:1060543.